



Results from the MAJORANA DEMONSTRATOR

Andrew Lopez
University of Tennessee Knoxville
On behalf of the MAJORANA Collaboration

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, the Particle Astrophysics and Nuclear Physics Programs of the National Science Foundation, and the Sanford Underground Research Facility.

Physics Motivation





Neutrino Mass Hierarchy Problem:

- Until recently neutrinos were thought to be massless
- The absolute neutrino mass scale is unknown
- Neutrino oscillation experiments can only measure the squared difference of the masses

Neutrinoless-Double Beta Decay:

- Hypothetical process in which only electrons are emitted
- Observable only if neutrinos are Majorana particles

If $0\nu\beta\beta$ decay is observed \Rightarrow
 Neutrinos are Majorana particles,
 Lepton number is violated,
 Sheds light on the absolute neutrino
 mass scale. ($\Gamma_{0\nu\beta\beta} \propto |m_{eff}|^2$)



Effective majorana mass as a function of the lightest neutrino mass


The MAJORANA DEMONSTRATOR



Operating 4850' underground at the Sanford Underground Research Facility, Lead, SD.

- Goals:**
- Demonstrate backgrounds low enough to justify building a tonne scale experiment.
 - Establish feasibility to construct & field modular arrays of Ge detectors.
 - Searches for additional physics beyond the standard model.


- Background Goal in the $0\nu\beta\beta$ peak region of interest (4 keV at 2039 keV)
3 counts/ROI-t-y (after analysis cuts); Measured Assay U.L. ≤ 3.5 counts/ROI-t-y
- Energy resolution of 2.4 keV FWHM @ 2039 keV (best of any $0\nu\beta\beta$ experiment)
- 44.1-kg of Ge detectors
 - 29.7 kg of 88% enriched ^{76}Ge crystals (35 detectors)
 - 14.4 kg of $^{\text{nat}}\text{Ge}$ (23 detectors)
 - Detector Technology: P-type, point-contact.
- 2 independent cryostats
 - ultra-clean, electroformed Cu
 - 22 kg of detectors per cryostat
 - naturally scalable
- Compact Shield
 - low-background passive Cu and Pb shield with active muon veto



N. Abgrall *et al.*, Adv. High Ener. Phys. **2014**, 365432 (2013)
arXiv:1308.1633

Funded by DOE Office of Nuclear Physics, NSF Particle Astrophysics, NSF Nuclear Physics with additional contributions from international collaborators.

8/3/2017



MAJORANA DEMONSTRATOR Implementation



Three Steps

Prototype cryostat: 7.0 kg (10) $^{\text{nat}}\text{Ge}$

Same design as Modules 1 and 2, but fabricated using OFHC Cu Components




Module 1: 16.9 kg (20) $^{\text{enr}}\text{Ge}$
5.6 kg (9) $^{\text{nat}}\text{Ge}$

Module 2: 12.9 kg (15) $^{\text{enr}}\text{Ge}$
8.8 kg (14) $^{\text{nat}}\text{Ge}$

June 2014-June 2015

9/2014: Module commissioning
5/2015 - 10/2015: In-shield running
10/2015 - 1/2016: Offline, upgrades
1/2016 - Present: in-shield running

4/2016: Module commissioning
7/2016 - Present: In-shield running



Advantages of ^{76}Ge detectors



- Intrinsic high-purity Ge detectors = source
- Excellent energy resolution:
approaching 0.1% at 2039 keV (~3 keV ROI)
- Demonstrated ability to enrich from 7.44% to $\geq 87\%$
- Powerful background rejection:
 - Granularity: multiple detectors hit
 - Pulse shape discrimination (PSD): multiple hits in a detector
 - Alpha events near surface: based on response





Ge Detector PSD efficiency



PSD cuts are optimized to keep 90% of single site and $< 10\%$ of multi-site events

- $0\nu\beta\beta$ is a single site event
- ^{208}Tl 2614 keV γ can pair produce and emit 2 γ , used to adjust PSD
- Both γ 's escape from detectors \rightarrow Double escape peak (DEP), single site
- One γ escapes from detectors \rightarrow Single escape peak (SEP), multi-site



^{208}Tl SEP (multiple site events) reduced to 6%

Delayed Charge Recovery and Alphas



Alpha background response observed in Module 1 commissioning (DS0)


- Identified as arising from alpha particles impinging on passivated surface

Results in prompt collection of some energy, plus very slow collection of remainder


Produces a distinctive waveform allowing a high efficiency cut

- “Delayed Charge Recovery” (DCR) parameter related to slope of tail

Example pole-zero corrected waveforms



Slow drift of charges along passivated surface results in very slow signal component



DCR paper arXiv:1610.03054



Detector Calibration



- Modules 1 and 2
- ^{228}Th calibration line source
- FWHM = 2.4keV at $Q_{\beta\beta}$ (2039 keV)

MaGe paper
Boswell et al. IEEE Trans.Nucl.Sci. 58 (2011)
[arXiv:1011.3827]

Calibration paper
arXiv:1702.02466




Comparison of a ^{228}Th line source simulation using MaGe and a measurement of M1. The simulated distribution was normalized by matching the integrals of both curves in the range from 2595 keV to 2635 keV.




DEMONSTRATOR Background Model

Radioassay paper: NIMA 828 (2016) 22 [arXiv:1601.03779]








Background Model paper
arXiv:1610.01146



Duty Cycles and Livetime



M1 Commissioning, no inner shield M1 inner shield M1 Multisampling Modules 1 and 2 Together in-shield Module 1 & 2 Integrated DAQ

	DS-0	DS-1	DS-2	DS-3	DS-4	DS-5
	Module 1	Module 1	Module 1	Module 1	Module 2	Module 1 & 2
	June 26 - Oct. 7, 2015	Dec. 31, 2015 - May 24, 2016	May 24 - July 14, 2016	Aug. 25 - Sep. 27, 2016	Aug. 25 - Sep. 27, 2016	Oct. 13, 2016 - May. 11 2017*
Total (days)	103.15	144.50	50.97	32.37	32.36	97.7
Total acquired	87.93	136.98	50.47	31.73	25.80	90.41
Physics  	47.70	61.34 + 20.41*	9.82 + 30.56*	29.97	23.84	82.52
High radon 	11.76	7.32	-	-	-	-
Calibration 	15.44	7.32	0.65	1.18	1.17	1.39
Down time 	15.21	7.51	0.50	0.64	6.56	7.29
Disruptive/Commissioning  	13.10	34.43 + 5.92*	2.41 + 7.03*	0.57	0.78	6.51

*Blind data


*Values up to Jan. 19, 2017

DS6 has started with multisampling and blindness.


$0\nu\beta\beta$ Region of Interest (DS-3 & DS-4)



- After cuts, 1 count in 400 keV window centered at 2039 keV ($0\nu\beta\beta$ peak)
 - Background index of 1.8×10^{-3} c/(keV kg y)
 - Projected background rate is $5.1^{+8.9}_{-3.2}$ c/(ROI t y) for a 2.9 keV (M1/DS3) & 2.6 keV (M2/DS4) keV ROI, (68%CL).
- Analysis cuts are still being optimized.






Muon Flux Measurement



- Measured total flux: $(5.31 \pm 0.17) \times 10^{-9} \mu/s/cm^2$.

Muon Flux paper
Astropart. Phys. 93, 70 (2017)
[arXiv:1602.07742]



Low Energy Program




- Low backgrounds and properties of the PPC HPGe detectors, allow for low energy searches for physics beyond the standard model.


Searches beyond SM

- Pseudoscaler dark matter
- Vector dark matter
- 14.4 keV Solar axion
- Pauli Exclusion Principle
- $e^- \rightarrow \nu\bar{\nu}\nu$

Low Energy paper
Phys. Rev. Lett. **118**, 161801 (2017)
[arXiv:1612.00886]



Pseudoscaler axion-like DM coupling



Vector particle DM coupling

MAJORANA DEMONSTRATOR Summary



- Commissioning is complete.
 - Both modules are collecting data in the final configuration.
- The ^{76}Ge enriched point contact detectors developed by MAJORANA
 - have attained the best energy resolution (2.4 keV FWHM at 2039 keV) of any $\beta\beta$ -decay experiment.
 - provide excellent pulse shape discrimination reduction of backgrounds.
 - at low energies have sub-keV energy thresholds and excellent resolution allowing the DEMONSTRATOR to perform sensitive test in this region for physics beyond the standard model.
- The DEMONSTRATOR's initial backgrounds are amongst the lowest backgrounds in the ROI achieved to date (approaching to GERDA's recent best value). Attained by development and selection of ultra-low activity materials and low mass designs.
- Combining the strengths of GERDA and the MAJORANA DEMONSTRATOR, the LEGEND Collaboration is moving forward with a ton-scale ^{76}Ge based experiment. Based on the successes to date, LEGEND should be able to reach the backgrounds ($\sim 0.1 \text{ c}/(\text{ROI t y})$) and energy resolution necessary for discovery level sensitivities in the inverted ordering region.

The MAJORANA COLLABORATION



The MAJORANA Collaboration



Black Hills State University, Spearfish, SD

Kara Keeter

Princeton University, Princeton, New Jersey

Graham K. Giovanetti

Duke University, Durham, North Carolina, and TUNL

Matthew Busch

Queen's University, Kingston, Canada

Ryan Martin

Joint Institute for Nuclear Research, Dubna, Russia

Viktor Brudanin, M. Shirchenko, Sergey Vasilyev, E. Yakushev, I. Zhitnikov

South Dakota School of Mines and Technology, Rapid City, South Dakota

Colter Dunagan, Cabot-Ann Christofferson, Anne-Marie Suriano, Jared Thompson

*Lawrence Berkeley National Laboratory, Berkeley, California and
the University of California - Berkeley*

Nicolas Abgrall, Yuen-Dat Chan, Lukas Hehn, Jordan Myslik, Alan Poon,
Kai Vetter

Tennessee Tech University, Cookeville, Tennessee

Mary Kidd

Technische Universität München, and Max Planck Institute, Munich, Germany

Tobias Bode, Susanne Mertens

Los Alamos National Laboratory, Los Alamos, New Mexico

Pinghan Chu, Steven Elliott, Ralph Massarczyk, Keith Rielage, Larry Rodriguez,
Harry Salazar, Brandon White, Brian Zhu

University of North Carolina, Chapel Hill, North Carolina, and TUNL

Thomas Caldwell, Thomas Gilliss, Chris Haufe, Reyco Henning, Mark Howe,
Samuel J. Meijer, Christopher O' Shaughnessy, Gulden Othman, Jamin Rager, Anna Reine,
Benjamin Shanks, Kris Vorren, John F. Wilkerson

*National Research Center 'Kurchatov Institute' Institute of Theoretical and
Experimental Physics, Moscow, Russia*

Alexander Barabash, Sergey Konovalov, Vladimir Yumatov

University of South Carolina, Columbia, South Carolina

Frank Avignone, Vince Guiseppe, David Tedeschi, Clint Wiseman

North Carolina State University, and TUNL

Matthew P. Green

University of South Dakota, Vermillion, South Dakota

CJ Barton, Wenqin Xu

Oak Ridge National Laboratory

Fred Bertrand, Charlie Havener, Monty Middlebrook, David Radford,
Robert Varner, Chang-Hong Yu

University of Tennessee, Knoxville, Tennessee

Yuri Efremenko, Andrew Lopez

Osaka University, Osaka, Japan

Hiroyasu Ejiri

University of Washington, Seattle, Washington

Sebastian Alvis, Tom Burritt, Micah Buuck, Clara Cuesta, Jason Detwiler, Julieta Gruszko,
Ian Guinn, David Peterson, Walter Pettus, R. G. Hamish Robertson, Nick Rouf,
Tim Van Wechel


Pacific Northwest National Laboratory, Richland, Washington

Isaac Arnquist, Eric Hoppe, Richard T. Kouzes




Backup Slides


Sensitivity, Background and Exposure



Discovery, Background and Exposure



Apparatus Overview



MAJORANA Underground Laboratory

